

The Open Access Israeli Journal of Aquaculture – Bamidgeh

As from **January 2010** The Israeli Journal of Aquaculture - Bamidgeh (IJA) will be published exclusively as **an on-line Open Access (OA)** quarterly accessible by all AquacultureHub (<http://www.aquaculturehub.org>) members and registered individuals and institutions. Please visit our website (<http://siamb.org.il>) for free registration form, further information and instructions.

This transformation from a subscription printed version to an on-line OA journal, aims at supporting the concept that scientific peer-reviewed publications should be made available to all, including those with limited resources. The OA IJA does not enforce author or subscription fees and will endeavor to obtain alternative sources of income to support this policy for as long as possible.

Editor-in-Chief

Dan Mires

Editorial Board

| | |
|-----------------------------|---|
| Rina Chakrabarti | Aqua Research Lab, Dept. of Zoology, University of Delhi, India |
| Angelo Colorni | National Center for Mariculture, IOLR Eilat, Israel |
| Daniel Golani | The Hebrew University of Jerusalem Jerusalem, Israel |
| Hillel Gordin | Kibbutz Yotveta, Arava, Israel |
| Sheenan Harpaz | Agricultural Research Organization Beit Dagan, |
| Gideon Hulata | Agricultural Research Organization Beit Dagan, |
| George Wm. Kissil | National Center for Mariculture, IOLR, Eilat, Israel |
| Ingrid Lupatsch | Swansea University, Singleton Park, Swansea, UK |
| Spencer Malecha | Dept. of Human Nutrition, Food & Animal Sciences, CTAHR, University of Hawaii |
| Constantinos Mylonas | Hellenic Center for Marine Research, Crete, Greece |
| Amos Tandler | National Center for Mariculture, IOLR Eilat, Israel |
| Emilio Tibaldi | Udine University Udine, Italy |
| Jaap van Rijn | Faculty of Agriculture, The Hebrew University of Jerusalem, Israel |
| Zvi Yaron | Dept. of Zoology, Tel Aviv University, Tel Aviv, Israel |

Published under auspices of
**The Society of Israeli Aquaculture and
Marine Biotechnology (SIAMB),
University of Hawai'i at Mānoa Library**

&

**University of Hawai'i at Mānoa
Aquaculture Program**
in association with
AquacultureHub

<http://www.aquaculturehub.org>



UNIVERSITY
of HAWAII
MĀNOA
LIBRARY



AquacultureHub.org

AquacultureHub
educate • learn • share • engage

ISSN 0792 - 156X

© Israeli Journal of Aquaculture - BAMIGDEH.

PUBLISHER:

Israeli Journal of Aquaculture - BAMIGDEH -
Kibbutz Ein Hamifratz, Mobile Post 25210,
ISRAEL

Phone: + 972 52 3965809


<http://siamb.org.il>

Copy Editor **Ellen Rosenberg**



The *IJA* appears exclusively as a peer-reviewed on-line open-access journal at <http://www.siamb.org.il>. To read papers free of charge, please register online at [registration form](#).

Sale of *IJA* papers is strictly forbidden.



An Arithmetic Index Based on Shell Height, Length, and Width, for Potential Selection of Soft-Body Wet Weight in Pacific Oyster, *Crassostrea gigas*

Xiaotong Wang^{1, *}, Ting Liu², Yongshan Liu¹, Peiyong Feng¹

¹ *School of Agriculture, Ludong University, Yantai 264025, China*

² *College of Life Science and Engineering, Shaanxi University of Science & Technology, Xi'an 710021, China*

(Received 27.12.13, Accepted 7.1.14)

Key words: Pacific oyster, selective breeding, soft-body wet weight, arithmetic product of shell length, height and width

Abstract

The Pacific oyster, one of the most widely grown bivalves, has a long commercial history, a large consumer base, and is one of the most important commercially cultured mollusks in the world. Only the soft body is edible therefore increasing the soft-body wet weight (SBWW) is the main breeding objective for oysters. The SBWW cannot be measured in-vivo. In general, the shell height (H) or shell length (L) are used to indirectly choose SBWW. But they are diverse and susceptible to environmental influences so they are not indicative of higher SBWW and inappropriate for the selection of optimal brooders. In this study, H, L as well as shell width (W) of wild and cultured populations were measured and the correlation coefficients between H, L, W, HLW (the arithmetic index based on shell height, length and width) and SBWW were calculated and found to be potential indicators for selective breeding of SBWW in Pacific oyster.

* Corresponding author. e-mail: wangxiaotong999@163.com

Introduction

Oysters have been cultured for well over a century. The Pacific oyster, *Crassostrea gigas*, is presently one of the most widely grown bivalves and one of the most important commercial mollusks around the world (FAO). Though some methods related to molecular breeding of oysters have been studied (Wang et al., 2011; Wang et al., 2012), conventional breeding including phenotypic selection are still the most efficient methods.

Increasing the SBWW is the main breeding objective of oyster culture, but it cannot be measured in-vivo. Generally, H or L are used to indirectly select SBWW. But there are huge differences in different environments (Mahon, 1983), and these indices are not appropriate for SBWW selection, therefore more accurate indices are required for this purpose.

Materials and Methods

Subjects. Two wild populations of pacific oysters from Penglai and Rushan and nine others cultured in Yantai (C1, C2, C3, C4, C5, C6, C7, C8, C9) in Shandong Province, China were used for this study.

Experimental methods. All oysters (11 groups, 30-78 each group) were brushed clean. Wild populations and cultured populations were temporarily held in plastic tanks for 3 days and fed with Spirulina powder.

The SBWW of each oyster was measured with an electronic scale. After weighing, each oyster was measured with Vernier calipers to determine shell height (maximum dorsal-ventral dimension), shell length (maximum anterior-posterior dimension), and shell width (maximum lateral dimension).

Data analysis. HLW was calculated using the following equation: $HLW = H * L * W$. The correlation coefficients between H, L, W, HLW, with SBWW were calculated and the significance of these correlation coefficients was also tested at the same time with CORR function in SAS (Statistics Analysis System) 9.0 software, respectively.

Results

The H, L, W and SBWW data was presented in supplementary materials (Table S1-11). As shown in Table 1 and Figure 1, in all populations, there are highly significant positive correlations between H, L, W, HLW, and SBWW ($p < 0.01$), but the correlation coefficients are different. The correlation coefficient between HLW and SBWW is the greatest (Red background) in each group.

In addition, the correlation coefficients between HLW and SBWW in cultured populations are greater than in wild populations (Figure 1 and Table 1).




Fig. 1. Comparison of the correlation coefficients between H, or L, or W, or HLW, and SBWW in different populations.

Table 1. The correlation coefficients and their significance level between H, or L, or W, or HLW, and SBWW.

| | Number of subjects | H-SBWW | L-SBWW | W-SBWW | HLW-SBWW |
|--------------|--------------------|---------|---------|---------|----------|
| Wild-Penglai | 53 | 0.54073 | 0.52786 | 0.58558 | 0.73393 |
| P-value | | <.0001 | <.0001 | <.0001 | <.0001 |
| Wild-Rushan | 41 | 0.53522 | 0.59047 | 0.34247 | 0.6625 |
| P-value | | 0.0003 | <.0001 | 0.0284 | <.0001 |
| Cultured_C1 | 32 | 0.57646 | 0.50184 | 0.60861 | 0.85457 |
| P-value | | 0.0006 | 0.0034 | 0.0002 | <.0001 |
| Cultured_C2 | 32 | 0.44572 | 0.80139 | 0.74882 | 0.83462 |
| P-value | | 0.0106 | <.0001 | <.0001 | <.0001 |
| Cultured_C3 | 30 | 0.56056 | 0.69381 | 0.66272 | 0.78059 |
| P-value | | 0.0013 | <.0001 | <.0001 | <.0001 |
| Cultured_C4 | 32 | 0.65992 | 0.8098 | 0.39319 | 0.87594 |
| P-value | | <.0001 | <.0001 | 0.026 | <.0001 |
| Cultured_C5 | 32 | 0.65992 | 0.8098 | 0.39319 | 0.87594 |
| P-value | | <.0001 | <.0001 | 0.026 | <.0001 |
| Cultured_C6 | 32 | 0.46364 | 0.77958 | 0.56687 | 0.82021 |
| P-value | | 0.0075 | <.0001 | 0.0007 | <.0001 |
| Cultured_C7 | 78 | 0.63955 | 0.66688 | 0.65319 | 0.76061 |
| P-value | | <.0001 | <.0001 | <.0001 | <.0001 |
| Cultured_C8 | 32 | 0.4917 | 0.56253 | 0.74192 | 0.86891 |
| P-value | | 0.0043 | 0.0008 | <.0001 | <.0001 |
| Cultured_C9 | 72 | 0.47825 | 0.55643 | 0.68021 | 0.74523 |
| P-value | | <.0001 | <.0001 | <.0001 | <.0001 |

Discussion

Table 1 and Figure 1 show that the correlation coefficient between HLW and SBWW in each population is greatest, suggesting that HLW is more suitable as the indirect index for SBWW selection than H, L, or W. Using HLW to select SBWW may promote breeding efficiency and accelerate breeding of Pacific oyster.

The correlation coefficients between HLW and SBWW in cultured populations are greater than in the two wild populations, indicating that a wild, varied, and complicated environment is likely to cause greater diversity in oyster shape.

Though many new selection methods are emerging (e.g. molecular marker assisted selection, MAS), phenotypic selection is still the basic and primary breeding method (Kingsolver et al., 2012; Piepho, et al., 2008), especially for mollusks because it is difficult to apply MAS to them. The arithmetic index based on shell height, length, and width (HLW) may be potentially useful for Pacific oyster breeding.

Acknowledgements

This research was supported by the National Natural Science Foundation of China (No. 31302181), Shandong Provincial Natural Science Foundation, China (No. ZR2013CM026), School Foundation of Ludong University (No. LY2013032) and Enterprise project (2013HX007).

References

FAO Fisheries & Aquaculture - Aquatic species.

Wang X., L. Li, F. Xu, and G. Zhang, 2011. Genomic DNA extraction from in vivo sampled tissue of Pacific oyster, *Crassostrea gigas*. *Isr. J. Aquacult. - Bamidgeh*, IIC:63.2011.671. 3 pages.

Wang X., X. Song, L. Li and G. Zhang, 2012. An improved method of extraction of DNA from the shell of Pacific oyster, *Crassostrea gigas*. *Isr. J. Aquacult. - Bamidgeh*, IJA:64.2012.800. 3 pages.

Mahon G.A.T., 1983. Selection goals in oyster breeding. *Aquaculture*, 33: 141-148.

Kingsolver J.G., S.E. Diamond, A.M. Siepielski, M.C. Stephanie, 2012. Synthetic analyses of phenotypic selection in natural populations: lessons, limitations and future directions. *Evol. Ecol.*, 26: 1101-1118.

Piepho H.P., J. Möhring, A.E. Melchinger, A. Büchse, 2008. BLUP for phenotypic selection in plant breeding and variety testing. *Euphytica*, 161: 209-228.

Supplementary materials (Table S1-11) can be downloaded at <http://dl.vmall.com/c00c28ar0a>.